

Table 5a. Reported Total Early Syphilis (TES)* Cases^ by Race and Select Demographics, Year to Date^

	Asian/Pacific Islander		Black		Hispanic		White		Other/Not Reported		Total	
<u>Sex</u>												
Female	0	0%	24	14%	0	0%	7	9%	0	0%	31	11%
Male	4	100%	143	85%	11	100%	69	90%	2	100%	229	88%
All	4	100%	167	100%	11	100%	76	100%	2	100%	260	100%
<u>Age at diagnosis</u>												
15 - 19	0	0%	17	10%	1	9%	1	1%	0	0%	19	7%
20 - 29	0	0%	80	47%	2	18%	16	21%	1	50%	99	38%
30 - 39	3	75%	33	19%	5	45%	15	19%	0	0%	56	21%
40 - 49	0	0%	28	16%	1	9%	27	35%	1	50%	57	21%
50+	1	25%	9	5%	2	18%	17	22%	0	0%	29	11%
All	4	100%	167	100%	11	100%	76	100%	2	100%	260	100%
<u>Region</u>												
CENTRAL	1	25%	56	33%	1	9%	30	39%	2	100%	90	34%
EASTERN	0	0%	76	45%	2	18%	15	19%	0	0%	93	35%
NORTHERN	3	75%	17	10%	8	72%	19	25%	0	0%	47	18%
NORTHWEST	0	0%	1	0%	0	0%	8	10%	0	0%	9	3%
SOUTHWEST	0	0%	17	10%	0	0%	4	5%	0	0%	21	8%
All	4	100%	167	100%	11	100%	76	100%	2	100%	260	100%

* Total Early Syphilis includes: Primary, Secondary and Early Latent Syphilis

^ See definitions under [Technical Notes](#)

Reported Cases of TES by Region, Year to Date

